

Schedule 1

Queensland fruit fly hosts and suspect hosts

Common Name	Scientific Name
Abiu	<i>Pouteria caimito</i>
Acerola	<i>Malpighia glabra</i>
Achachairu	<i>Garcinia humilis</i>
Apple	<i>Malus domestica</i>
Apricot, Cherry, nectarine, peach, peacharine, plum, plumcot	<i>Prunus</i> spp.
Avocado	<i>Persea americana</i>
Babaco	<i>Carica pentagona</i>
Banana	<i>Musa acuminata</i>
Black sapote	<i>Diospyros ebenum</i>
Blueberry	<i>Vaccinium corymbosum</i>
Blackberry, boysenberry, loganberry, youngberry, raspberry	<i>Rubus</i> spp.
Brazil cherry	<i>Eugenia uniflora</i>
Breadfruit	<i>Artocarpus altilis</i>
Caimito (star apple)	<i>Chrysophyllum cainito</i>
Cape gooseberry	<i>Physalis peruviana</i>
Capsicum	<i>Capsicum annum</i>
Carambola	<i>Averrhoa carambola</i>
Cashew apple	<i>Anacardium occidentale</i>
Casimiroa (white sapote)	<i>Casimiroa edulis</i>
Cherimoya, custard apple, soursop, sweetsop	<i>Annona</i> spp.
Chilli	<i>Capsicum annum</i> <i>Capsicum frutescens</i>
Chinese mulberry*	<i>Cudrania tricuspidata</i>
Citron, finger lime, grapefruit, lemon, lime,	<i>Citrus</i> spp.

mandarin, Orange, pummelo, tangelo	
Climbing fig*	<i>Ficus pumila</i>
Clivia*	<i>Clivia</i> spp.
Coffee berry	<i>Coffea</i> spp.
Date (fresh)	<i>Phoenix dactylifera</i>
Durian	<i>Durio zibethinus</i>
Eggplant	<i>Solanum melongena</i>
Feijoa	<i>Feijoa sellowiana</i>
Fig	<i>Ficus carica</i>
Granadilla	<i>Passiflora quadrangularis</i>
Grapes	<i>Vitis</i> spp.
Grumichama	<i>Eugenia braziliensis</i>
Guava	<i>Psidium</i> spp.
Hog plum	<i>Spondias mombin</i>
Indian hawthorn*	<i>Rhaphiolepis indica</i>
Jaboticaba	<i>Myrciaria cauliflora</i>
Jackfruit	<i>Artocarpus heterophyllus</i>
Jew plum	<i>Spondias dulcis</i>
Kiwifruit	<i>Actinidia deliciosa</i>
Kumquat	<i>Fortunella japonica</i>
Lilly pilly*	<i>Syzygium smithii</i>
Logan	<i>Dimocarpus longan</i>
Loofah	<i>Luffa acutangula</i>
Loquat	<i>Eriobotrya japonica</i>
Lychee	<i>Litchii chinensis</i>
Mango	<i>Mangifera indica</i>
Mangosteen	<i>Garcinia mangostana</i>
Miraclefruit	<i>Synsepalum dulcificum</i>
Moreton fig*	<i>Ficus macrophylla</i>
Mulberry	<i>Morus nigra</i>
Nashi	<i>Pyrus pyrifolia</i> var. <i>culta</i>
Olive	<i>Olea europaea</i>
Ornamental pear*	<i>Pyrus calleryana</i>
Passionfruit	<i>Passiflora</i> spp.
Pawpaw	<i>Carica papaya</i>
Pear	<i>Pyrus communis</i>
Pepino	<i>Solanum muricatum</i>
Persimmon	<i>Diospyros kaki</i>
Pomegranate	<i>Punica granatum</i>
Prickly pear	<i>Opuntia stricta</i> ; <i>O. ficus-indica</i>

Quince	<i>Cydonia oblonga</i>
Rambutan	<i>Nephelium lappaceum</i>
Rollinia	<i>Rollinia deliciosa</i>
Rose apple	<i>Syzygium jambos</i>
Santol	<i>Sandoricum indicum</i>
Sapodilla	<i>Manilkara zapota</i>
Sapote	<i>Casimiroa edulis</i>
Strawberry	<i>Fragaria spp.</i>
Tamarillo	<i>Solanum bataceum</i>

Tomato	<i>Lycopersicon esculentum</i>
Wax jambu/ water apple	<i>Syzygium samarangense</i>

*indicates suspect host

Dr Michelle Rodan
ACTING EXECUTIVE DIRECTOR
BIOSECURITY

Date: 27 January 2021